



urological surgical associates

Jerold Grubman MD, FACS
Andrei Kachala MD, FACS
Benjamin Fand MD, FACS
William Terens MD, FACS
Joshua L. Wein MD, FACS
Rupa Patel MD

10 Parsonage Road, Suite 117, Edison, N.J. 08837 Tel: (732) 494 9400 Fax: (732) 548 3931
3 Hospital Plaza, Suite 200, Old Bridge, N.J. 08857 Tel: (732) 679 2010 Fax: (732) 679 2077

General Dietary Recommendations for the Prevention of Stones

1. **Increase overall fluid intake.** Goal should be drink enough fluids to urinate more than 2 liters of urine per day.
2. **Decrease protein intake.** Excess protein intake can contribute to both uric acid and calcium based stones. Limit intake to 6-8 ounces of beef, pork, poultry or fish per day.
3. **Decrease salt intake.** Excess salt intake can cause extra calcium to enter into the urine and thus lead to stone formation.
4. **Increase "Citrate" intake.** Citrate is commonly found in lemons. Adding lemon juice to your fluids or drinking fresh lemonade can help inhibit the formation of stones
5. **Consume adequate calcium.** Contrary to popular belief stone formers should NOT restrict their calcium intake. It is encouraged that you eat 2 servings of calcium rich foods daily.
6. **Decrease Oxalate Intake.** If you form calcium oxalate stones it is essential you limit your oxalate intake. Oxalate containing foods include spinach, rhubarb, beets, chocolate, peanuts and teas. For a more comprehensive list of foods please refer to our low oxalate diet form.